



# Alterations of local functional connectivity in lifespan: A resting-state fMRI study

Xin Wen<sup>1</sup> | Hui He<sup>2</sup> | Li Dong<sup>2,3</sup> | Junjie Chen<sup>1</sup> | Jie Yang<sup>3</sup> | Hao Guo<sup>1</sup> | Cheng Luo<sup>2,3</sup> | Dezhong Yao<sup>2,3</sup>

<sup>1</sup>College of Information and Computer, Taiyuan University of Technology, Taiyuan, China

<sup>2</sup>The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for Neuroinformation, University of Electronic Science and Technology of China, Chengdu, China

<sup>3</sup>School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, China

## Correspondence

Li Dong, The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for Neuroinformation, University of Electronic Science and Technology of China, Chengdu, China.  
Email: lidong@uestc.edu.cn

Junjie Chen, College of Information and Computer, Taiyuan University of Technology, Taiyuan, Shanxi, China.  
Email: chenjj@tyut.edu.cn

## Funding information

This study was financially supported by the National Natural Science Foundation of China (Nos. 81861128001, 81701778, 61672374, 61876124) and the Sichuan Science and Technology Program (2018JZ0073, 2019YJ0179).

## Abstract

**Introduction:** As aging attracted attention globally, revealing changes in brain function across the lifespan was largely concerned. In this study, we aimed to reveal the changes of functional networks of the brain (via local functional connectivity, local FC) in lifespan and explore the mechanism underlying them.

**Materials and Methods:** A total of 523 healthy participants (258 males and 265 females) aged 18–88 years from part of the Cambridge Center for Ageing and Neuroscience (CamCAN) were involved in this study. Next, two data-driven measures of local FC, local functional connectivity density (lFCD) and four-dimensional spatial-temporal consistency of local neural activity (FOCA), were calculated, and then, general linear models were used to assess the changes of them in lifespan.

**Results:** Local functional connectivity (lFCD and FOCA) within visual networks (VN), sensorimotor network (SMN), and default mode network (DMN) decreased across the lifespan, while within basal ganglia network (BGN), local connectivity was increased across the lifespan. And, the fluid intelligence decreased within BGN while increased within VN, SMN, and DMN.

**Conclusion:** These results might suggest that the decline of executive control and intrinsic cognitive ability in the aging population was related to the decline of functional connectivity in VN, SMN, and DMN. Meanwhile, BGN might play a regulatory role in the aging process to compensate for the dysfunction of other functional systems. Our findings may provide important neuroimaging evidence for exploring the brain functional mechanism in lifespan.

## KEYWORDS

four-dimensional spatial-temporal consistency of local neural activity, lifespan, local functional connectivity, local functional connectivity density, resting-state fMRI

Wen and He contributed equally to this study.

All the three institutions contributed equally to this study.

The peer review history for this article is available at <https://publons.com/publon/10.1002/brb3.1652>

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Brain and Behavior* published by Wiley Periodicals LLC.

## 1 | INTRODUCTION

With the intensification of global aging, lifespan research has become a hot spot (Bookheimer et al., 2019; Taylor et al., 2017; Wei, Zhuang, et al., 2018). Recently, the development and application of functional networks via functional connectivity (FC) have provided new perspectives and discoveries for changes in brain function across the lifespan (Han et al., 2018; Nazeri et al., 2015; Vij, Nomi, Dajani, & Uddin, 2018). In previous behavioral studies (Grady, 2012; Mansson et al., 2015; Smith et al., 2015), it was suggested that with the development of the human brain (young-mature-aging), some cognitions and behaviors including motor ability, vision, and auditory would exhibit an inverted U-shaped curve. Such a phenomenon could be explained that cognition and behavior showed a trend of increasing first and then decreasing with age (Damoiseaux, 2017; Wing et al., 2018). Using functional magnetic resonance imaging (fMRI), several brain function researches indicated that the elderly have lower functional connectivity within the default mode network (DMN), dorsal attention network (DAN), sensorimotor network (SMN), visual (VN) and fronto-parietal networks (FPN; Betzel et al., 2014; Cassady et al., 2019; Ferreira & Busatto, 2013; Grady, Sarraf, Saverino, & Campbell, 2016; Spreng, Stevens, Viviano, & Schacter, 2016), as well as higher functional connectivity within the basal ganglia network (BGN; Allen, 2011). Meanwhile, structural MRI-related studies have shown that in addition to temporal lobe-related networks, older people have lower gray matter volumes in the DMN, DAN, and auditory networks (Damoiseaux, 2017; Liu, 2017). In addition, several previous lifespan and aging studies showed that older people had higher functional connectivity between networks (such as DMN, FPN, DAN, and cingular-opercular network [CON]; Damoiseaux, 2017; Grady et al., 2016; Wang, Su, Shen, & Hu, 2012; Xia et al., 2019). In general, as the aging process deepens, the functional connectivity within network decreased and the functional connections between networks increased (Ferreira & Busatto, 2013; Vij et al., 2018). Reduced functional connectivity may be due to reduced cognitive and other functions, and the increase in functional connectivity between networks may be due to the compensatory mechanism of the functional network (Damoiseaux, 2017; Ferreira & Busatto, 2013; Naik, Banerjee, Bapi, Deco, & Roy, 2017; Vij et al., 2018). Furthermore, BGN is associated with a variety of functions, including motor, cognitive, motivational, and emotional processes (Doya, 2000), and has overlapping several functions with other networks, especially with SMN which could be expected that BGN may compensate to the dysfunction of SMN (Figley, 2017; Yanagisawa, 2018). Meanwhile, Regners et al. also found that increased effective connectivity from dDMN to BGN in long-term abstinence may be a compensatory mechanism related to behavioral monitoring (Castellazzi, 2014; Regner et al., 2016). However, the physiological mechanisms of functional network changes in lifespan are still unclear, especially whether and how the potential compensation mechanism is in lifespan.

In recent years, resting-state fMRI has been used as a powerful tool for exploring spontaneous brain activity and changes in

brain function (Biswal, Zerrin Yetkin, Haughton, & Hyde, 1995; De Luca, Beckmann, De Stefano, Matthews, & Smith, 2006). Unlike traditional task fMRI focusing on a single functional system at a time, resting-state fMRI can provide important spontaneous activity information of functional connectivity for the interpretation of lifespan functional network changes (Betzel et al., 2014; Li et al., 2019; Tomasi & Volkow, 2012; Yang, 2016). Considering that the local functional homogeneity of spontaneous activity is neurobiologically relevant due to possible anatomical, developmental, and neurocognitive factors (Jiang et al., 2015; Wei, Chang, et al., 2018), there are two popular resting-state measures of local functional connectivity (local FC). One is the local functional connectivity density (IFCD), which reflects the local functional hubs of the brain in the time domain (Tomasi & Volkow, 2010), and another is the four-dimensional spatial-temporal consistency of local neural activity (FOCA), which reflects the functional state of the brain locally in the time-frequency domain (Azeez & Biswal, 2017; Dong et al., 2015). These two local FC measures may comprehensively assess the local functional activity, more comprehensively, from its specific perspective. Therefore, with the advantages of satisfactory reproducibility and reliability, both data-driven IFCD and FOCA measures of local FC may be suitable for revealing changes in brain function and its mechanism across the lifespan.

This study was the first to explore the alterations of functional networks in lifespan and the possible underlying mechanism through local FCs of IFCD and FOCA, to our knowledge. In this study, we hypothesized that the compensation across the lifespan may exist in a wide range of high-level (e.g., DMN) and primary networks (e.g., VN and SMN) via local functional connectivity, and BGN may play important role in the compensatory mechanism. Using resting-state fMRI, IFCD and FOCA were calculated on healthy participants aged 18–88 years. Then, a general linear model (GLM) was used to assess the relationships between local functional indicators and age/age<sup>2</sup>, while controlling the nuisance items of the gender, head motion, and intracranial volume. In addition, relationships between functional measures and behavior scores were also investigated using GLM, while adding the nuisance items as covariates. We predicted that it may provide important neuroimaging evidence for exploring the brain mechanism in lifespan.

## 2 | MATERIALS AND METHODS

### 2.1 | Participants

In this study, 523 healthy participants (258 males and 265 females) aged 18–88 years from part of the Cambridge Center for Ageing and Neuroscience (CamCAN, <http://www.cam-can.org/>; Taylor et al., 2017) were involved. Among all the participants, the proportion of male and female participants remained equally, and there were about 100 participants per 10-year-old. Participants performed cognitive tasks outside the MRI scanner. The tests used

in this study were the Cattell Culture Fair Test (complete nonverbal puzzles involving series completion, classification, matrices, and conditions), to assess fluid intelligence (Horn & Cattell, 1966), and the speed choice reaction time (RT) task (participants were required to respond as quickly as possible to 1 of 4 possible cued fingers using a 4-button response box, total trial was 67), to assess speed of processing. For the RT tasks, the mean (M-RT) and variability (SD of RT values, SD-RT) were computed from individual trials. Moreover, due to the lack of records and data quality control, some participants were excluded in this study (see Table 1 and Table S2 for details). All procedures followed the Helsinki Declaration and have been approved by the Cambridgeshire 2 Research Ethics Committee, a local ethics committee in the UK.

## 2.2 | fMRI data acquisition and preprocessing

Imaging data were collected by a 3T MRI scanner (Siemens TIM Trio). During the scanning period, all participants were required to lie still and keep their eyes closed. The T1-weighted anatomical images were gathered using an MPRAGE sequence with the following parameters: TR/TE = 2,250 ms/2.99 ms; flip angle = 9°; FOV = 256×240 × 192 mm<sup>3</sup>; voxel size = 1 × 1 × 1 mm<sup>3</sup>. For resting-state fMRI measurements, 261 volumes of echo-planar imaging (EPI) sequences were acquired with the following parameters: sequential descending order; slice thickness 3.7 mm with a slice gap of 20% for whole-brain coverage; TR/TE = 1,970 ms/30 ms; flip angle = 78°; FOV = 192×192 mm<sup>2</sup>; voxel size = 3×3 × 4.44 mm<sup>3</sup>; number of slices = 32; duration time = 520 s.

The resting-state fMRI data were preprocessed using SPM12 as implemented in the Neuroscience Information Toolbox (NIT, version 1.3, <http://www.neuro.uestc.edu.cn/NIT.html>; Dong, 2018). The first 5 volumes were deleted from the resting-state fMRI data of each subject for excluding T1 saturation effects. Then, the images were preprocessed including realignment, slice time correction, spatial normalization using T1-weighted MRI data (3 × 3 × 3 mm<sup>3</sup>). The head motion of each participant was calculated using the mean framewise displacement (mean FD; Power, Barnes, Snyder, Schlaggar, & Petersen, 2012). Participants whose FD was two or more SD above the group mean FD were excluded from further analysis.

**TABLE 1** Information of participants

	Number	Male/ female	Range (years/ values)	Mean	SD
Age	523	258/265	18–88	51.28	17.60
Cattell	509	254/255	12–44	32.82	6.37
RT_M	476	236/240	0.35–1.11	0.57	0.13
RT_SD	476	236/240	0.04–0.39	0.12	0.06


## 2.3 | IFCD and FOCA

In this work, IFCD and FOCA measures were calculated to assess the local brain activity in resting state using NIT software (<http://www.neuro.uestc.edu.cn/NIT.html>, version 1.3; Dong, 2018). The local functional connectivity density (IFCD) reflects the distribution of functional hubs of the human brain from the perspective of time consistency in the local region (Tomasi & Volkow, 2010). For a voxel  $v$  in the brain, the IFCD value of voxel  $v$  was the number of voxels in the contiguous functional connectivity cluster in a local region (Tomasi & Volkow, 2010). The Pearson correlation was used to assess the strength of the functional connectivity between voxel  $v$  and its closest neighbors of voxels functionally connected, and a given time courses threshold of 0.6 and the MRI signal-to-noise ratio threshold of 50 were used to ensure significant functional connectivity (Tomasi & Volkow, 2010). A voxel was added to the list of voxels connected with voxel  $v$  only if it was linked to voxel  $v$  by a continuous path of connected voxels and correlation was larger than the given threshold and this computation repeated for the next voxel in the list until no new neighbors can be added to the list. This calculation was performed for all voxels, and the individual map was obtained by dividing by the mean value of the map. Before calculating IFCD maps, normalized functional images were passband-filtered (0.01–0.08 Hz) and nuisance signals, including 12 head motion parameters (6 parameters of translation and rotation and their derivative), linear trend, global mean, individual mean white matter, and cerebrospinal fluid signals, were removed from the unsmoothed fMRI data.

On the other hand, the four-dimensional spatial-temporal consistency of local neural activity (FOCA) reflects the functional state of the brain locally consist in time–frequency domain (the consistency from both temporal homogeneity of local adjacent voxels using temporal correlation and regional stability of brain activity states between neighboring time points; Azeez & Biswal, 2017; Dong et al., 2015). For each voxel, FOCA value was calculated as mean temporal and spatial correlation of a local region from normalized functional images. After calculating FOCA of all voxels across the brain, the FOCA value was divided by the mean value of the whole brain. Brain regions with high FOCA values are considered to have higher consistency of local spontaneous activity. Before calculating FOCA maps, nuisance signals, including 12 head motion parameters, linear trend, individual mean white matter, and cerebrospinal fluid signals, were removed from the unsmoothed fMRI data. The detailed information of FOCA can be seen in Dong et al. (2015). Finally, the IFCD image and FOCA image were smoothed with an 8 mm FWHM Gaussian kernel.

## 2.4 | Statistical analysis

To exhibit the relationship between local functional connectivity and age, we used GLM to fit local measures (IFCD and FOCA maps)



**FIGURE 1** Relationships between local functional connectivity (FOCA and IFCD) and age/age<sup>2</sup>. (a) Impact of age on local FC and (b) impact of age<sup>2</sup> on local FC (cluster size > 30,  $p < .05$ , false discovery rate (FDR) correction). Red areas indicate that local FC increased significantly as age increased, while blue areas indicate the opposite. R stands for the right hemisphere of the brain, and L stands for the left hemisphere of the brain

over age/age<sup>2</sup> while adding the gender ( $X_{sex}$ ), head motion (mean framewise displacement,  $X_{mFD}$ ), and intracranial volume ( $X_{volume}$ ) as covariates (Equation 1). Additionally, GLM was also conducted to fit linear local measures over behavior scores (fluid intelligence and speed choice reaction time) adding the same covariates ( $X_{behavior}$  in Equation 2). Because the changes of mean FCs across lifespan were mainly linear and quadratic (the inverted U shape; Grady, 2012) and FC changes within some functional networks (VN, CEN) have linear decreases across lifespan while those within DMN have quadratic (the inverted U shape) decreases, and SMN, SAN shown both linear and quadratic decreases of FCs (Vij et al., 2018; Wang et al., 2012). It is reasonable to apply a GLM to reveal the age effect on local FCs. Furthermore, the age and behavioral score were strongly correlated ( $r^2 = .43$ ,  $p = 3.6 \times 10^{-63}$ , see Figure S1). Because adding the behavioral score in the GLM in Equation 1 would lead to a collinearity problem, according to the previous studies (Kievit, 2014; Onoda, Ishihara, & Yamaguchi, 2012), two GLMs were applied to these two measures, respectively.

$$\text{localFC} = \beta_0 + \beta_1 \times \text{age} + \beta_2 \times \text{age}^2 + \beta_3 \times X_{sex} + \beta_4 \times X_{mFD} + \beta_5 \times X_{volume} \quad (1)$$

$$\text{localFC} = \beta'_0 + \beta'_1 \times X_{behavior} + \beta'_2 \times X_{sex} + \beta'_3 \times X_{mFD} + \beta'_4 \times X_{volume} \quad (2)$$

For each local functional measure (IFCD and FOCA maps), the T value of coefficients of age and age<sup>2</sup> in Equation 1 ( $\beta_1$  and  $\beta_2$ , respectively) was used to measure the impact of age (linear and quadratic) on local functional measures. Meanwhile, the T value of coefficients of behavior score (Cattell score, M\_RT, and SD\_RT) in Equation 2 ( $\beta'_1$ ) was used to measure the significance of the linear relationship between them and local functional measures. All the significance level was set to  $p < .05$ , false discovery rate (FDR)-corrected.

### 3 | RESULTS

#### 3.1 | Changes of IFCD and FOCA

First, we analyzed the linear correlation between local functionality (IFCD and FOCA) and age. During the resting state, the IFCD values

**TABLE 2** Detailed information on each brain region for the relationship between local FC and age

Index	Region	MNI coordinates			Peak t_value	Cluster size		
		X	Y	Z				
Age								
IFCD	Olfactory_L	-27	12	-18	8.34	9,204		
	Temporal_Pole_Sup_L	-45	9	-12	8.17			
	Temporal_Sup_R	42	6	-12	8.09			
	Insula_R	27	18	-18	7.26			
	Hippocampus_R	24	-30	-3	7.15			
	Amygdala_R	21	3	-12	6.92			
	Temporal_Sup_L	-45	-9	0	6.83			
	Olfactory_R	18	6	-15	6.63			
	Hippocampus_L	-24	-24	-6	6.57			
	Cerebellum_9_L	-9	-45	-42	6.57			
	Cerebellum_9_R	3	-48	-42	6.23			
	Caudate_R	9	18	12	6.17			
	Occipital_Mid_R	15	-93	6	-7.52		4,248	
	Calcarine_L	-21	-96	3	-7.13			
	Occipital_Mid_L	-27	-93	0	-6.95			
	Cuneus_R	6	-90	27	-6.48			
	Calcarine_R	3	-75	12	-6.27			
	Lingual_L	-9	-72	-3	-5.57			
	frontal_Inf_Tri_R	39	21	27	-4.92			736
	Precentral_L	-48	6	30	-4.56			434
Frontal_Sup_Medial_L	0	18	42	-4.62	208			
FOCA	Thalamus_L	-18	-24	0	9.32	6,903		
	Thalamus_R	21	-21	0	9.57			
	Caudate_L	-9	15	15	8.8			
	Hippocampus_R	24	-30	-3	8.38			
	Caudate_R	9	15	15	8.34			
	Hippocampus_L	-33	-36	0	7.56			
	Temporal_Pole_Sup_L	-45	12	-16	7.23			
	Frontal_Mid_L	-30	60	30	4.36		139	
	Frontal_Sup_Medial_L	0	57	45	4		172	
	Calcarine_R	3	-66	21	-5.94		7,491	
	Paracentral_Lobule_R	3	-33	51	-5.72			
	Lingual_R	9	-75	-3	-5.49			
	Cerebellum_6_L	-27	-78	-18	-5.35			
	Cerebellum_Crus1_L	-45	-54	-24	-5.35			
	Cingulum_Mid_R	3	-15	45	-5.08			
	Postcentral_R	42	-27	48	-4.68			
	Postcentral_L	-45	-27	45	-3.54		902	
	Frontal_Mid_Orb_L	0	48	-12	-5.65		258	
	Cerebellum_Crus1_R	42	-60	-24	-4.98		231	

in the olfactory cortex, the superior temporal gyrus, right insula, hippocampus, right amygdala, cerebellum inferior, and right caudate nucleus were positively correlated with age, while areas that negatively correlated with IFCD involved occipital gyrus, calcarine, left cuneus, left lingual gyrus, inferior frontal gyrus, left precentral gyrus, and left

medial superior frontal gyrus. At the same time, the FOCA values of thalamus, caudate nucleus, hippocampus, superior temporal gyrus, left middle frontal gyrus, and left medial superior frontal gyrus were positively correlated with age; the areas that negatively correlated with FOCA involved right calcarine, right paracentral lobule, right

lingual gyrus, cerebellum superior, cerebellum superior, right medial and lateral cingulate gyrus, postcentral gyrus, and left superior orbital frontal gyrus (Figure 1a, detailed information located in Table 2). Meanwhile, results of IFCD with time course thresholding of 0.5 and 0.7 were similar to the above findings and are shown in Figures S2 and S3; and there also were wide changes of degree of global networks at the voxel level in the brain across lifespan (Figure S4).

In addition, a nonlinear correlation between local functionality (IFCD and FOCA) and age was analyzed. The IFCD values in the right dorsolateral superior frontal gyrus, left hippocampus, left precentral gyrus, left precuneus, right inferior temporal gyrus, and right caudate nucleus were positively correlated with age, while area that negatively correlated with IFCD involved right postcentral gyrus. At the same time, the FOCA values of the inferior temporal gyrus, caudate nucleus, and right inferior frontal orbital gyrus were positively correlated with age; the areas that negatively correlated with FOCA involved cerebellum superior, right calcarine, precentral gyrus, superior temporal gyrus, left postcentral gyrus, and transverse temporal gyrus (Figure 1b, detailed information located in Table 3). In addition, gender and age together have few impacts on local FCs in our study (Figure S5).

### 3.2 | Relationship between behavior and local FCs

In order to illustrate the relationship between behavior and local FCs, we calculated the correlation between them. The fluid intelligence (Cattell) was negatively correlated with age in the lifespan

while positively related to local FC within DMN, VN and partial SMN. However, there were few significant relationships between local FCs and RT\_M/RT\_SD (Figure 2). The detailed information could be seen in Table S1.

## 4 | DISCUSSION


This study explored the alterations in functional networks across the lifespan and the underlying mechanism via local functional connectivity. In brief, as shown in Figure 3, it was found that local functional connectivity (IFCD and FOCA) in the VN, SMN, and DMN decreased across the lifespan while that in the BGN increased across the lifespan. Such results might be explained by the decline in the VN, SMN, and DMN, reflecting the impairment of corresponding functions, while the increase in the BGN indicates compensation for functional networks.

### 4.1 | Changes of IFCD and FOCA across the lifespan

The changes of functional connectivity in lifespan have always been a research hot spot, and there have been a few studies on the mechanism behind the changes of functional connectivity. Previous studies have shown that functional connectivity in lifespan decreased with age, mainly within DMN, FPN, CON, SMN, and VN, while some studies suggested that FCs within the partial motor and


Index	Region	MNI coordinates			Peak t_value	Cluster size
		X	Y	Z		
Age <sup>2</sup>						
IFCD	Frontal_Sup_R	21	12	57	5.91	190
	Hippocampus_L	-36	-9	-24	4.92	367
	Precentral_L	-30	-3	51	4.6	118
	Precuneus_L	-9	-54	51	4.31	71
	Temporal_Inf_R	45	-6	-39	4.03	58
	Caudate_R	12	18	12	4	46
	Postcentral_R	39	-21	48	-3.77	41
FOCA	Temporal_Inf_R	48	-45	-6	5.96	78
	Caudate_R	15	21	15	5.42	197
	Caudate_L	-18	9	24	4.18	152
	Temporal_Inf_L	-57	-27	-21	4.12	510
	Frontal_Inf_Orb_R	42	36	-12	3.97	67
	Cerebellum_6_L	-24	-81	-18	-4.57	908
	Calcarine_R	6	-81	3	-4.11	
	Precentral_L	-42	-12	57	-4.65	77
	Temporal_Sup_L	-57	0	0	-4.03	113
	Postcentral_L	-54	-15	42	-4.03	66
	Heschl_R	54	-12	9	-3.98	59
	Temporal_Sup_R	60	0	0	-3.78	

**TABLE 3** Detailed information on each brain region for the relationship between local FC and age<sup>2</sup>



**FIGURE 2** Relationships between local functional connectivity (FOCA and IFCD) and behavior scores (cluster size > 30 voxels; Cattell-fluid intelligence,  $p < .05$ , FDR correction; RT\_M, RT\_SD-average response time in response time tasks, intraindividual response time variability in response time tasks,  $p < .001$ , uncorrected). Red areas indicate that local FC decreased significantly as behavior scores decreased, while blue areas indicate the opposite. R stands for the right hemisphere of the brain, and L stands for the left hemisphere of the brain

**FIGURE 3** Illustration of the underlying mechanisms of four networks over the lifespan. As age increases, the local FCs within the VN, SMN, and DMN decreased (impairment), while those within the BGN increased (compensation). The behavior scores exhibited an inverted U-shaped curve across the lifespan



subcortical networks increased with age (Damoiseaux, 2017; Ferreira & Busatto, 2013). In general, the decline of situational memory, self-reference processing, and thinking wandering in the elderly leads to the decline of functional connectivity in DMN (Buckner, Andrews-Hanna, & Schacter, 2008). Moreover, the functional connectivity between DMN and FPN was positively correlated with age, which may be due to the weakening of the separation of the two functional networks with aging (Grady et al., 2016; Spreng et al., 2016). On the other hand, Coste and his colleagues found that the decline

in functional connectivity within CON may be due to decreased alertness among older people (Coste & Kleinschmidt, 2016) and Yan, Zhuo, Wang, and Wang (2011) suggested that the decline of VN internal functional connectivity with age may be due to the decline of visual function in the elderly. In addition, the elderly performed an impaired motor performance, yet the consistent changes of FC were not observed within SMN. As the age grows, FC increased within some portions of SMN, while some portions of SMN exhibited the contrary phenomena. The lifespan and aging studies implied that

the declines within SMN were associated with decreases in sensory processing and diminished communications within a “sensorimotor feedback” system (Roski, 2013; Seidler et al., 2015). Moreover, fluid intelligence is associated with executive control and intrinsic cognitive ability, which decreases with age growing indicating the elderly might impair with the above abilities (Finn et al., 2015; Gray, Chabris, & Braver, 2003). Our study found that local functional connectivity (IFCD and FOCA) within DMN, VN, and SMN decreased in lifespan. We also found the fluid intelligence was negatively correlated with age in the lifespan while positively related to local FC within DMN, VN, and partial SMN. Together, our results indicate that the decline of fluid intelligence might be linked with decreased local functional connectivity within VN, SMN, and DMN.

#### 4.2 | The possible compensatory role of the BGN

It was widely believed that the increased FC changes in lifespan could reflect a compensatory mechanism, and most of these changes were intranetwork FC. For example, the FCs between DMN and FPN/attentional networks were increased in lifespan due to the dedifferentiation process of aging brain, occurrence of aging for late neurodevelopmental stages, and a compensatory for cognitive decline (Grady et al., 2016; Zhai & Li, 2019; Zonneveld et al., 2019). Meanwhile, the BGN was thought to be related to motor learning functions and the increased FC within BGN might suggest as a compensatory role of dysfunctions of motor learning, especially in Parkinson's disease-related research (Lyman, Anguera, & Terman, 2011; Saling & Phillips, 2008). Furthermore, it was also found that the homologous basal ganglia circuitry was related to cognition and emotion (Saling & Phillips, 2008) and an increased FC within it might compensate high-order cognition (Siman-Tov, 2016). In lifespan, the BGN might play a compensatory role to balance the dysfunction of other functional networks. In our study, the local FC (IFCD and FOCA) with BGN increased in lifespan reflects that the increased FCs in BGN might provide the compensatory for the decline function networks of primary and high-order cognition. In summary, the BGN might regulate other functional networks (VN, SMN, and DMN) and compensate the corresponding cognitive decline in lifespan.

#### 4.3 | Limitations

Although our findings could give insight into the compensatory role of the BGN across the lifespan to some degree, several limitations should be addressed. First, behavioral data were relatively insufficient. Further research should involve more behavioral data to reveal more detailed cognitive relationships between local FCs, such as working memory, motor ability, and emotion recognition. Second, fMRI has low temporal resolution compared to electroencephalogram (EEG). In the future, EEG data should be involved to study real-time effective behavior results. Finally, because of the accurate compensatory effects are not tested in this work, the accurate

mechanism of the BGN remains unclear. Our study perhaps offers one possible explanation of the compensatory mechanism of the BGN across the lifespan through local FCs, and more research efforts are needed in this regard. Additionally, since the distribution of functional hubs in the human brain is a critical topic, it would be helpful to expand our research with complex network analysis in future work.

## 5 | CONCLUSION

In this study, we used resting-state IFCD and FOCA measures to investigate the changes of local functional connectivity in lifespan and tried to give the underlying mechanism: The decline of cognitive function in the aging population led to the decline of FC in VN, SMN, and DMN. Meanwhile, BGN may play a regulatory role in the aging process to compensate for the dysfunction of other functional systems.

#### ACKNOWLEDGMENT

We wish to thank the Cambridge Center for Ageing and Neuroscience for providing the data publicly.

#### CONFLICT OF INTEREST

All authors have no conflicts of interest to disclose.

#### AUTHOR CONTRIBUTIONS

LD and JC conceived the project. XW, HH, and LD implemented the statistical analysis. JY preprocessed the data. XW, HH, and LD wrote and revised the paper. HH, DY, HG, and CL provided critical suggestions for the manuscript.

#### DATA AVAILABILITY STATEMENT

Publicly available datasets were analyzed in this study. This data can be found here: <https://camcanarchive.mrc-cbu.cam.ac.uk/dataaccess/>.

#### ORCID

Xin Wen  <https://orcid.org/0000-0003-2363-1190>

Cheng Luo  <https://orcid.org/0000-0003-0524-5886>

#### REFERENCES

- Allen, E. A., Erhardt, E. B., Damaraju, E., Gruner, W., Segall, J. M., Silva, R. F., ... Calhoun, V. D. (2011). A baseline for the multivariate comparison of resting-state networks. *Frontiers in Systems Neuroscience*, 5, 2.
- Azeez, A. K., & Biswal, B. B. (2017). A Review of resting-state analysis methods. *Neuroimaging Clinics of North America*, 27(4), 581–592. <https://doi.org/10.1016/j.nic.2017.06.001>
- Betzel, R. F., Byrge, L., He, Y., Goñi, J., Zuo, X.-N., & Sporns, O. (2014). Changes in structural and functional connectivity among resting-state networks across the human lifespan. *NeuroImage*, 102(Pt 2), 345–357. <https://doi.org/10.1016/j.neuroimage.2014.07.067>
- Biswal, B. B., Zerrin Yetkin, F., Haughton, V. M., & Hyde, J. S. (1995). Functional connectivity in the motor cortex of resting human brain



- using echo-planar MRI. *Magnetic Resonance in Medicine*, 34(4), 537–541. <https://doi.org/10.1002/mrm.1910340409>
- Bookheimer, S. Y., Salat, D. H., Terpstra, M., Ances, B. M., Barch, D. M., Buckner, R. L., ... Yacoub, E. (2019). The Lifespan Human Connectome Project in Aging: An overview. *NeuroImage*, 185, 335–348. <https://doi.org/10.1016/j.neuroimage.2018.10.009>
- Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: Anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences*, 1124, 1–38.
- Cassady, K., Gagnon, H., Lalwani, P., Simmonite, M., Foerster, B., Park, D., ... Polk, T. A. (2019). Sensorimotor network segregation declines with age and is linked to GABA and to sensorimotor performance. *NeuroImage*, 186, 234–244. <https://doi.org/10.1016/j.neuroimage.2018.11.008>
- Castellazzi, G., Palesi, F., Casali, S., Vitali, P., Sinforiani, E., Wheeler-Kingshott, C. A., & D'Angelo, E. (2014). A comprehensive assessment of resting state networks: Bidirectional modification of functional integrity in cerebro-cerebellar networks in dementia. *Frontiers in Neuroscience*, 8, 223.
- Coste, C. P., & Kleinschmidt, A. (2016). Cingulo-opercular network activity maintains alertness. *NeuroImage*, 128, 264–272. <https://doi.org/10.1016/j.neuroimage.2016.01.026>
- Damoiseaux, J. S. (2017). Effects of aging on functional and structural brain connectivity. *NeuroImage*, 160, 32–40. <https://doi.org/10.1016/j.neuroimage.2017.01.077>
- De Luca, M., Beckmann, C. F., De Stefano, N., Matthews, P. M., & Smith, S. M. (2006). fMRI resting state networks define distinct modes of long-distance interactions in the human brain. *NeuroImage*, 29(4), 1359–1367. <https://doi.org/10.1016/j.neuroimage.2005.08.035>
- Dong, L., Luo, C., Cao, W., Zhang, R., Gong, J., Gong, D., & Yao, D. (2015). Spatiotemporal consistency of local neural activities: A new imaging measure for functional MRI data. *Journal of Magnetic Resonance Imaging*, 42(3), 729–736. <https://doi.org/10.1002/jmri.24831>
- Dong, L., Luo, C., Liu, X., Jiang, S., Li, F., Feng, H., ... Yao, D. (2018). Neuroscience information toolbox: An open source toolbox for EEG-fMRI multimodal fusion analysis. *Frontiers in Neuroinformatics*, 12, 56.
- Doya, K. (2000). Complementary roles of basal ganglia and cerebellum in learning and motor control. *Current Opinion in Neurobiology*, 10(6), 732–739. [https://doi.org/10.1016/S0959-4388\(00\)00153-7](https://doi.org/10.1016/S0959-4388(00)00153-7)
- Ferreira, L. K., & Busatto, G. F. (2013). Resting-state functional connectivity in normal brain aging. *Neuroscience and Biobehavioral Reviews*, 37(3), 384–400. <https://doi.org/10.1016/j.neubiorev.2013.01.017>
- Figley, T. D., Moghadam, M., Bhullar, N., Kornelsen, J., Courtney, S. M., & Figley, C. R. (2017). Probabilistic white matter atlases of human auditory, Basal Ganglia, Language, precuneus, sensorimotor, visual and visuospatial networks. *Frontiers in Human Neuroscience*, 11, 306.
- Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., ... Constable, R. T. (2015). Functional connectome fingerprinting: Identifying individuals using patterns of brain connectivity. *Nature Neuroscience*, 18(11), 1664–1671. <https://doi.org/10.1038/nn.4135>
- Grady, C. (2012). The cognitive neuroscience of ageing. *Nature Reviews Neuroscience*, 13(7), 491–505. <https://doi.org/10.1038/nrn3256>
- Grady, C., Sarraf, S., Saverino, C., & Campbell, K. (2016). Age differences in the functional interactions among the default, frontoparietal control, and dorsal attention networks. *Neurobiology of Aging*, 41, 159–172. <https://doi.org/10.1016/j.neurobiolaging.2016.02.020>
- Gray, J. R., Chabris, C. F., & Braver, T. S. (2003). Neural mechanisms of general fluid intelligence. *Nature Neuroscience*, 6(3), 316–322. <https://doi.org/10.1038/nn1014>
- Han, L., Savalia, N. K., Chan, M. Y., Agres, P. F., Nair, A. S., & Wig, G. S. (2018). Functional parcellation of the cerebral cortex across the human adult lifespan. *Cerebral Cortex*, 28(12), 4403–4423. <https://doi.org/10.1093/cercor/bhy218>
- Horn, J. L., & Cattell, R. B. (1966). Refinement and test of the theory of fluid and crystallized general intelligences. *Journal of Educational Psychology*, 57(5), 253–270. <https://doi.org/10.1037/h0023816>
- Jiang, L., Xu, T., He, Y., Hou, X.-H., Wang, J., Cao, X.-Y., ... Zuo, X.-N. (2015). Toward neurobiological characterization of functional homogeneity in the human cortex: Regional variation, morphological association and functional covariance network organization. *Brain Structure and Function*, 220(5), 2485–2507. <https://doi.org/10.1007/s00429-014-0795-8>
- Kievit, R. A., Davis, S. W., Mitchell, D. J., Taylor, J. R., Duncan, J., Henson, R. N., & Cam-CAN Research Team (2014). Distinct aspects of frontal lobe structure mediate age-related differences in fluid intelligence and multitasking. *Nature Communications*, 5, 5658.
- Li, H., Cao, W., Zhang, X., Sun, B., Jiang, S., Li, J., ... Luo, C. (2019). BOLD-fMRI reveals the association between renal oxygenation and functional connectivity in the aging brain. *NeuroImage*, 186, 510–517. <https://doi.org/10.1016/j.neuroimage.2018.11.030>
- Liu, K., Yao, S., Chen, K., Zhang, J., Yao, L., Li, K., ... Guo, X. (2017). Structural brain network changes across the adult lifespan. *Frontiers in Aging Neuroscience*, 9, 275.
- Lyman, K., Anguera, J., & Terman, D. (2011). A model of healthy aging and motor inhibition in the basal ganglia. *BMC Neuroscience*, 12, 331. <https://doi.org/10.1186/1471-2202-12-S1-P331>
- Mansson, K. N., Frick, A., Boraxbekk, C.-J., Marquand, A. F., Williams, S. C. R., Carlbring, P., ... Furmark, T. (2015). Predicting long-term outcome of Internet-delivered cognitive behavior therapy for social anxiety disorder using fMRI and support vector machine learning. *Translational Psychiatry*, 5, e530. <https://doi.org/10.1038/tp.2015.22>
- Naik, S., Banerjee, A., Bapi, R. S., Deco, G., & Roy, D. (2017). Metastability in Senescence. *Trends in Cognitive Sciences*, 21(7), 509–521. <https://doi.org/10.1016/j.tics.2017.04.007>
- Nazeri, A., Chakravarty, M. M., Rotenberg, D. J., Rajji, T. K., Rath, Y., Michailovich, O. V., & Voineskos, A. N. (2015). Functional consequences of neurite orientation dispersion and density in humans across the adult lifespan. *Journal of Neuroscience*, 35(4), 1753–1762. <https://doi.org/10.1523/JNEUROSCI.3979-14.2015>
- Onoda, K., Ishihara, M., & Yamaguchi, S. (2012). Decreased functional connectivity by aging is associated with cognitive decline. *Journal of Cognitive Neuroscience*, 24(11), 2186–2198. [https://doi.org/10.1162/jocn\\_a\\_00269](https://doi.org/10.1162/jocn_a_00269)
- Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage*, 59(3), 2142–2154. <https://doi.org/10.1016/j.neuroimage.2011.10.018>
- Regner, M. F., Saenz, N., Maharajh, K., Yamamoto, D. J., Mohl, B., Wylie, K., ... Tanabe, J. (2016). Top-down network effective connectivity in abstinent substance dependent individuals. *PLoS ONE*, 11(10), e0164818. <https://doi.org/10.1371/journal.pone.0164818>
- Roski, C., Caspers, S., Langner, R., Laird, A. R., Fox, P. T., Zilles, K., ... Eickhoff, S. B. (2013). Adult age-dependent differences in resting-state connectivity within and between visual-attention and sensorimotor networks. *Frontiers in Aging Neuroscience*, 5, 67.
- Saling, L. L., & Phillips, J. G. (2008). Age-related slowing of movement as basal ganglia dysfunction. *European Review of Aging and Physical Activity*, 5(2), 69–77. <https://doi.org/10.1007/s11556-008-0036-3>
- Seidler, R., Erdeniz, B., Koppelmans, V., Hirsiger, S., Méritat, S., & Jäncke, L. (2015). Associations between age, motor function, and resting state sensorimotor network connectivity in healthy older adults. *NeuroImage*, 108, 47–59. <https://doi.org/10.1016/j.neuroimage.2014.12.023>
- Siman-Tov, T., Bosak, N., Sprecher, E., Paz, R., Eran, A., Aharon-Peretz, J., & Kahn, I. (2016). Early age-related functional connectivity decline in high-order cognitive networks. *Frontiers in Aging Neuroscience*, 8, 330.

- Smith, S. M., Nichols, T. E., Vidaurre, D., Winkler, A. M., Behrens, T. E. J., Glasser, M. F., ... Miller, K. L. (2015). A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nature Neuroscience*, *18*(11), 1565–1567. <https://doi.org/10.1038/nn.4125>
- Spreng, R. N., Stevens, W. D., Viviano, J. D., & Schacter, D. L. (2016). Attenuated anticorrelation between the default and dorsal attention networks with aging: Evidence from task and rest. *Neurobiology of Aging*, *45*, 149–160. <https://doi.org/10.1016/j.neurobiolaging.2016.05.020>
- Taylor, J. R., Williams, N., Cusack, R., Auer, T., Shafto, M. A., Dixon, M., ... Henson, R. N. (2017). The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) data repository: Structural and functional MRI, MEG, and cognitive data from a cross-sectional adult lifespan sample. *NeuroImage*, *144*(Pt B), 262–269. <https://doi.org/10.1016/j.neuroimage.2015.09.018>
- Tomasi, D., & Volkow, N. D. (2010). Functional connectivity density mapping. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(21), 9885–9890. <https://doi.org/10.1073/pnas.1001414107>
- Tomasi, D., & Volkow, N. D. (2012). Aging and functional brain networks. *Molecular Psychiatry*, *17*(5), 549–558. <https://doi.org/10.1038/mp.2011.81>
- Vij, S. G., Nomi, J. S., Dajani, D. R., & Uddin, L. Q. (2018). Evolution of spatial and temporal features of functional brain networks across the lifespan. *NeuroImage*, *173*, 498–508. <https://doi.org/10.1016/j.neuroimage.2018.02.066>
- Wang, L., Su, L., Shen, H., & Hu, D. (2012). Decoding lifespan changes of the human brain using resting-state functional connectivity MRI. *PLoS ONE*, *7*(8), e44530. <https://doi.org/10.1371/journal.pone.0044530>
- Wei, D., Zhuang, K., Ai, L., Chen, Q., Yang, W., Liu, W., ... Qiu, J. (2018). Structural and functional brain scans from the cross-sectional Southwest University adult lifespan dataset. *Scientific Data*, *5*, 180134.
- Wei, Y., Chang, M., Womer, F. Y., Zhou, Q., Yin, Z., Wei, S., ... Wang, F. (2018). Local functional connectivity alterations in schizophrenia, bipolar disorder, and major depressive disorder. *Journal of Affective Disorders*, *236*, 266–273. <https://doi.org/10.1016/j.jad.2018.04.069>
- Wing, E. A., Iyengar, V., Hess, T. M., LaBar, K. S., Huettel, S. A., & Cabeza, R. (2018). Neural mechanisms underlying subsequent memory for personal beliefs: An fMRI study. *Cognitive, Affective, & Behavioural Neuroscience*, *18*(2), 216–231. <https://doi.org/10.3758/s13415-018-0563-y>
- Xia, Y., Chen, Q., Shi, L., Li, M. Z., Gong, W., Chen, H., & Qiu, J. (2019). Tracking the dynamic functional connectivity structure of the human brain across the adult lifespan. *Human Brain Mapping*, *40*(3), 717–728. <https://doi.org/10.1002/hbm.24385>
- Yan, L., Zhuo, Y., Wang, B., & Wang, D. J. J. (2011). Loss of coherence of low frequency fluctuations of BOLD FMRI in Visual Cortex Of Healthy Aged Subjects. *Open Neuroimaging Journal*, *5*(5), 105–111. <https://doi.org/10.2174/18744440001105010105>
- Yanagisawa, N. (2018). Functions and dysfunctions of the basal ganglia in humans. *Proceedings of the Japan Academy, Series B*, *94*(7), 275–304. <https://doi.org/10.2183/pjab.94.019>
- Yang, A. C., Tsai, S. J., Liu, M. E., Huang, C. C., & Lin, C. P. (2016). The association of aging with white matter integrity and functional connectivity hubs. *Frontiers in Aging Neuroscience*, *8*, 143.
- Zhai, J., & Li, K. (2019). Predicting brain age based on spatial and temporal features of human brain functional networks. *Frontiers in Human Neuroscience*, *13*, 62.
- Zonneveld, H. I., Pruim, R. H. R., Bos, D., Vrooman, H. A., Muetzel, R. L., Hofman, A., ... Vernooij, M. W. (2019). Patterns of functional connectivity in an aging population: The Rotterdam Study. *NeuroImage*, *189*, 432–444. <https://doi.org/10.1016/j.neuroimage.2019.01.041>

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Wen X, He H, Dong L, et al. Alterations of local functional connectivity in lifespan: A resting-state fMRI study. *Brain Behav.* 2020;10:e01652. <https://doi.org/10.1002/brb3.1652>